



UNIVERSITÀ
DEGLI STUDI
FIRENZE

FLORE

Repository istituzionale dell'Università degli Studi di Firenze

Low Florbetapir PET Uptake and Normal A β 1-42 Cerebrospinal Fluid in an APP Ala713Thr Mutation Carrier

Questa è la Versione finale referata (Post print/Accepted manuscript) della seguente pubblicazione:

Original Citation:

Low Florbetapir PET Uptake and Normal A β 1-42 Cerebrospinal Fluid in an APP Ala713Thr Mutation Carrier / Lombardi, Gemma; Berti, Valentina; Tedde, Andrea; Bagnoli, Silvia; Piaceri, Irene; Polito, Cristina; Lucidi, Giulia; Ferrari, Camilla; Ginestroni, Andrea; Moretti, Marco; Pupi, Alberto; Nacmias, Benedetta; Sorbi, Sandro. - In: JOURNAL OF ALZHEIMER'S DISEASE. - ISSN 1387-2877. - ELETTRONICO. - 57:(2017), pp. 697-703. [10.3233/JAD-161170]

Availability:

This version is available at: 2158/1082872 since: 2017-05-16T09:08:10Z

Published version:

DOI: 10.3233/JAD-161170

Terms of use:

Open Access

La pubblicazione è resa disponibile sotto le norme e i termini della licenza di deposito, secondo quanto stabilito dalla Policy per l'accesso aperto dell'Università degli Studi di Firenze (<https://www.sba.unifi.it/upload/policy-oa-2016-1.pdf>)

Publisher copyright claim:

(Article begins on next page)

Gemma Lombardi^a, Valentina Berti^a, Andrea Tedde^a, Silvia Bagnoli^a, Irene Placenta^a, Cristina Polito^b, Giulia Lucidi^{a,c}, Camilla Ferrari^c, Andrea Ginestroni^d, Marco Morini^a, Alberto Pupi^b, Benedetta Nacmias^{a,*} and Sandro Sorbi^{a,c}

^a*Department of Neuroscience, Psychology, Drug Research and Child Health, University of Florence, Florence, Italy*

^b*Department of Biomedical, Experimental and Clinical Sciences “Mario Serio”, Nuclear Medicine, University of Florence, Florence, Italy*

^c*IRCCS Don Gnocchi, Florence, Italy*

^d*Neuroradiology Unit AOU Careggi, Florence, Italy*

Handling Associate Editor: Daniela Galimberti

Accepted 31 January 2017

Abstract. According to the literature, the *APP* Ala713Thr mutation is associated with Alzheimer’s amyloid angiopathy. We describe a case of dementia clinically compatible with frontotemporal dementia in a *APP* Ala713Thr mutation carrier in which both [¹⁸F]Florbetapir PET uptake and Aβ₁₋₄₂ cerebrospinal fluid levels were low. Further evidences are required to establish if this association is only incidental.

Keywords: *APP* Ala713Thr, familial Alzheimer’s disease, florbetapir PET, frontotemporal dementia

INTRODUCTION

The revised National Institute on Aging-Alzheimer’s Association (NIA-AA) diagnostic criteria for dementia due to Alzheimer’s disease (AD) encourage the use of biomarkers in order to improve AD cases identification [1], and according

to International Working Group-2 diagnostic criteria, the diagnosis of AD requires the presence of a pathophysiological biomarker. The accuracy of amyloid positron emission tomography (PET) imaging in detecting amyloid plaques has been demonstrated [3], and several studies have found a correlation between amyloid PET and CSF fluid (CSF) Aβ₁₋₄₂ levels and cognitive decline. Amyloid plaques in the brain [4] as well as amyloid PET uptake [5]. In the context of AD causative mutation, the certainty that AD is caused by AD pathology is in-

*Correspondence to: Prof. Benedetta Nacmias, Department of Neuroscience, Psychology, Drug Research and Child Health, University of Florence, Viale Pieraccini 6, 50139 Florence, Italy. Tel.: +39 055 7948910; Fax: +39 0552758265; E-mail: benedetta.nacmias@unifi.it.

A 51-year-old Italian man was admitted to our clinic because of gradual change in behavior and personality, cognitive decline, and urinary incontinence; symptoms started one year before the first visit. Family history included a maternal aunt affected by unspecified early onset dementia (55 years) who died at the age of 75. No other cases of neurodegenerative/vascular dementia were known in the family. The mother died at 76 years because of renal failure in diabetes mellitus, while the father was affected by syphilis and died at the age of 78 years.

Information about health status of his cousins were not available. The patient underwent clinical-neuropsychological assessment and conventional brain magnetic resonance imaging (MRI); subsequently, since the diagnosis was uncertain, supplemental investigations on biomarkers were performed: brain [^{18}F]fluorodeoxyglucose ([^{18}F]FDG) PET, CSF analysis, genetic analysis for frontotemporal dementia (FTD), genetic analysis for FAD and lastly amyloid PET. All exams, except the amyloid PET, were carried out one year after the symptoms onset (baseline); neuropsychological evaluation was performed at baseline and at 6-month follow-up (Table 1), while amyloid PET was made 3 years after the symptoms onset.

The extensive neuropsychological evaluation was conducted with the Mini-Mental State Examination (MMSE) [10] and by tests assessing memory (Rey Auditory Verbal Learning Test [11], Babcock Story Recall Test [12], Digit Span Forward and Backward [13]), attention (Trail Making Test A and B [12]), language (Verbal Fluency [11, 14]), executive functions (Frontal Assessment Battery [15], Stroop Test [16]), visuospatial abilities (Rey–Osterrieth Complex Figure Test [17]), praxia (qualitative evaluation), mood

assessment.

With regards to the genetic analysis, DNA was extracted from peripheral blood cells by standard procedures, by utilizing the QIAzol lysis reagent (Qiagen, Hilden, Germany). PCR for FTD causative mutations (*MAPT*, *MAPT* H1 and *MAPT* H2) and for FAD causative mutations (*APP*) were performed following the manufacturer's protocol [23]. The Apolipoprotein E genotype was determined using PCR and HinfI digestion. A combination of the two single nucleotide polymorphisms (rs429358 and rs7412) on the *APOE* gene was used. Lumbar puncture and CSF collection were performed according to good clinical practice. The sample was analyzed for total-Tau, phosphorylated-Tau (p-Tau), $\text{A}\beta_{1-42}$ by commercial enzyme-linked immunosorbent assay (innogenetics ELISA). The reference values were: for total-Tau <300 pg/ml, for p-Tau <60 pg/ml, and for $\text{A}\beta_{1-42}$ >60 pg/ml.

RESULTS

At neurological examination, signs of mild cognitive impairment and left hand resting tremor without other extrapyramidal signs. Neuropsychological assessment showed significant apathy, reduced verbal initiative and verbal fluency, short-term memory impairment, slight executive functions, and functional decline (instrumental self-care deficit). The MMSE score was below reference norms (21), the Neuropsychiatric Inventory score was relatively high (28) because of delusions, severe apathy, motor disinhibition, typed behaviors, dietary changes, and daytime sleepiness. The clinical picture was suggestive of early onset dementia in the mild spectrum.

Trail Making Test		
A	(55'')	(50'')
B	73'' (110'')/1 Err	not evaluable
Stroop Test		
Time Interference Effect	17.5'' (18)	not evaluable
Errors Interference Effect	(0)	not evaluable
Phonemic Word Fluency	6.5 (5)	11.5 (10)
Semantic Word Fluency	20.5 (19)	14.5 (13)
Neuropsychiatric Inventory	(28/144)	(40/144)
Activities of Daily Living Scale	(4/6 preserved)	(3/6 preserved)

Brain MRI showed diffuse fronto-temporo-parietal atrophy (Pasquier score was 14 and the most affected regions were the fronto-temporals) with bilateral involvement of hippocampus (Scheltens score was 3 on the right side, 2 on the left side), without evident vascular burden (Fazekas score was 1) (Fig. 1a). The MRI pattern was not suggestive for any specific dementia subtype.

Brain [^{18}F]FDG PET scan (Fig. 1b) revealed severe bilateral hypometabolism in the anterior prefrontal, anterior cingulate, medial frontal, orbitofrontal cortices, in the hippocampi, amygdala, temporal pole, and caudate nuclei (mainly on the right side). Moderate bilateral hypometabolism in the insula and posterior cingulate cortex was also evident, whereas the metabolism in the posterolateral parietal cortex was preserved. This pattern of predominantly anterior hypometabolism has been interpreted as suggestive of neurodegenerative disease not in the AD spectrum but more likely in the FTD spectrum.

With regards to CSF biomarkers, $\text{A}\beta_{1-42}$ and phospho-Tau were within the normal range (respectively, 771 pg/ml and 35 pg/ml), whereas total-Tau was slightly increased (354 pg/ml); according to current evidence, these values and their ratios were not suggestive of AD (total Tau/ $\text{A}\beta_{1-42}$ value (0.46) was not suggestive of AD profile according to Duits

et al. [25], phospho-Tau/ $\text{A}\beta_{1-42}$ not suggestive of AD profile according to Duits et al. [26]).

Based on these results and on the clinical diagnosis of behavioral FTD was made.

Genetic analysis for FTD causative mutations (*MAPT*, *GRN*, *C9orf72*) was negative. In the absence of early onset dementia, genetic analysis for causative mutations was also performed. The occurrence of an *APP* Ala713T mutation (heterozygous mutation in the presence of a wild-type genotype).

Lastly, amyloid imaging with [^{18}F]AV-45 PET was performed and no significant amyloid retention was found (Fig. 1c). The late disease onset, the clinical picture characterized by mutism and compulsive behavior, the presence of attention and executive dysfunction, gait, and dysautonomic dysfunction (e.g., hypersalivation).

According to all the data, an atypical form of AD, the so-called "frontal variant AD", in the absence of a causative mutation and in the absence of CSF biomarkers, has been assumed. According to the non-genetic biomarkers profile, according to Rascovsky's criteria [27], a clinical picture of possible FTD, behavioral variant of AD, was postulated.




Fig. 1. Neuroimaging data a) MRI, axial T2-weighted fluid-attenuated inversion recovery images and coronal T2-weighted images, evidence of cortical and hippocampal atrophy; b) [^{18}F]FDG-PET: severe hypometabolism in frontal and caudate nuclei; c) [^{18}F]Florbetapir PET: no tracer retention in grey matter.

DISCUSSION

The potential AD pathogenicity of *APP* Ala713Thr (g.275329G>A) variant is confirmed by segregation of the mutation with the disease [8, 9, 28, 29], absence of the mutation in control subjects [8, 9, 28], and results of bioinformatics analysis [30].

This variant is reported in mutation databases as pathogenic (<http://www.molgen.ua.ac.be/ADmutations> [31], <http://www.alzforum.org/mutations> [32], <http://www.hgmd.cf.ac.uk/> [33]); it has been described associated with autosomal dominant FAD, with both early and late onset, and cerebrovascular lesions. Despite these considerations, asymptomatic subjects of heterozygous carriers have been reported [8, 9]; since these cases were mainly younger than 65 years [8] or than the average age at onset of respective affected family members [9], they cannot be considered spared from the disease. Moreover, an

incomplete penetrance of the mutation was proposed in order to explain this phenotype. Genetic or environmental factors could be responsible for the expression/unexpression of the mutation [7].

In the present case, the genetic diagnosis contrasts with the lack of evidence of disease as assessed by both CSF analysis and neuroimaging. Some cases of negative amyloid PET have been reported in FAD in both Arctic APP mutations [34] and *APP* Glu693del mutation [35], also in rare sporadic cases of disease. The absence of compound B PET ([^{11}C]PiB PET) and the expected cerebral amyloid burden.

PiB binding was found to be specific for certain forms of $\text{A}\beta$, such as $\text{A}\beta_{42}$ and $\text{A}\beta_{40}$, or high-affinity binding sites within the plaques present in each amyloid plaque. The absence of [^{18}F]Florbetapir uptake in the

variant (oligomeric toxicity without alteration in A β ratio [37]), it is conceivable that CSF A β ₁₋₄₂ value remains in the normal range and that CSF A β ₁₋₄₀ (not evaluated in this case) decreases. This hypothesis is based on the evidence that the reduced A β -peptides in CSF match the peptides accumulated in the amyloid plaques [38].

Instead of AD diagnosis, the co-occurrence of an altered A β sequence with a FTD phenotype could suggest an association between *APP* Ala713Thr mutation and the FTD pathology. A causative role of *APP* Ala713Thr in pathology different from AD has not been previously supposed, whereas mutation located at the same codon (the *APP* Ala713Val) was found associated to schizophrenia with cognitive deficit [39], suggesting that mutations involving codon 713 can be pleiotropic.

In conclusion, we describe the first case of *APP* Ala713Thr mutation carrier associated to both low [¹⁸F]Florbetapir PET uptake and normal A β ₁₋₄₂ CSF. It remains to be clarified whether *APP* Ala713Thr mutation is linked to AD without *in vivo* evidence of A β biomarkers or more likely to other pathology as FTD, and the debate is unsolvable without histopathological examination. Based on our findings, we suggest that in the occurrence of an AD causative mutation, more than one criterion is necessary for the *in vivo* evidence of AD pathology and that pathophysiological biomarkers should be studied even in carriers of a pathogenic mutation, in order to confirm the genetic diagnosis or, alternatively, to suppose other diagnosis and the relative pathogenic mechanism.

ACKNOWLEDGMENTS

This study was supported by grants from Italian Ministry of Health RF-2010-2319722 and

- [2] Dubois B, Feldman HH, Jacova C, et al (2007) Prevalence of Alzheimer's disease in a population-based autopsy study. *Neurology* 68, 205-209.
- [3] Dubois B, Feldman HH, Jacova C, et al (2007) Prevalence of Alzheimer's disease in a population-based autopsy study. *Neurology* 68, 205-209.
- [4] Strozyk D, Blennow K, White LR, et al (2005) Abeta 42 levels correlate with cognitive decline in a population-based autopsy study. *Neurology* 65, 652-656.
- [5] Fagan AM, Mintun MA, Machuga CT, et al (2007) CS, Shah AR, LaRossa GN, Spiro AJ, Mathis CA, DeKosky ST, Morris JC (2007) Inverse relation between *in vivo* amyloid and cerebrospinal fluid Abeta42 in humans. *Neurology* 68, 512-519.
- [6] Dubois B, Feldman HH, Jacova C, et al (2007) Prevalence of Alzheimer's disease in a population-based autopsy study. *Neurology* 68, 205-209.
- [7] Carter DA, Desmarais E, Bellis M, et al (1992) More missense in amyloid gene. *Neurology* 42, 256.
- [8] Rossi G, Giaccone G, Maletta R, et al (2007) Mangieri M, Giovagnoli AR, Bizzini C, Di Natale M, Tagliavini F, Bugiani M (2007) A family with Alzheimer disease with A713T mutation of the *APP* gene. *Neurology* 68, 910-912.
- [9] Conidi ME, Bernardi L, Puccio G, et al (2015) Curcio SA, Colao R, Piscopo P, Galliani F, Clodomiro A, Mirabelli M, Torchia G, Di Lorenzo R, Mandich RG, Bruni AC (2015) Homozygous mutation in an autosomal dominant form of Alzheimer's disease. *Neurology* 84, 2266-2273.
- [10] Measso G, Cavarzeran F, Zappalà G, et al (1993) TH, Pirozzolo FJ, Amaducci LA, et al (1993) Prevalence of Alzheimer's disease in a population-based autopsy study. *Neurology* 41, 205-209.

- [15] Appollonio I, Leone M, Isella V, Piamarta F, Consoli T, Villa ML, Forapani E, Russo A, Nichelli P (2005) The Frontal Assessment Battery (FAB): Normative values in an Italian population sample. *Neurol Sci* **26**, 108-116.
- [16] Caffarra P, Vezzadini G, Dieci F, Zonato A, Venneri A (2002) Una versione abbreviata del test di Stroop: Dati normativi nella popolazione italiana. *Nuova Riv Neurol* **12**, 111-115.
- [17] Caffarra P, Vezzadini G, Dieci F, Zonato F, Venneri A (2002) Rey-Osterrieth complex figure: Normative values in an Italian population sample. *Neurol Sci* **22**, 443-447.
- [18] Kaufer DI, Cummings JL, Ketchel P, Smith V, MacMillan A, Shelley T, Lopez OL, DeKosky ST (2000) Validation of the NPI-Q, a brief clinical form of the Neuropsychiatric Inventory. *J Neuropsychiatry Clin Neurosci* **12**, 233-239.
- [19] Katz S, Ford AB, Moskowitz RW, Jackson BA, Jaffe MW (1963) Studies of illness in the aged. The index of ADL: A standardized measure of biological and psychosocial function. *JAMA* **185**, 914-919.
- [20] Scheltens P, Barkhof F, Leys D, Pruvo JP, Nauta JJ, Vermersch P, Steinhilber M, Valk J (1992) Atrophy of medial temporal lobes on MRI in "probable" Alzheimer's disease and normal ageing: Diagnostic value and neuropsychological correlates. *J Neurol Neurosurg Psychiatry* **55**, 967-972.
- [21] Pasquier F, Leys D, Weerts JGE, Mounier-Vehier F, Barkhof F, Scheltens P (1996) Inter- and intraobserver reproducibility of cerebral atrophy assessment on MRI scans with hemispheric infarcts. *Eur Neurol* **36**, 268-272.
- [22] Fazekas F, Chawluk JB, Alavi A, Hurtung HI, Zimmerman RA (1987) MR signal abnormalities at 1.5 T in Alzheimer's dementia and normal aging. *Am J Roentgenol* **149**, 351-356.
- [23] Nacmias B, Piaceri I, Bagnoli S, Tedde A, Piacentini S, Sorbi S (2014) Genetics of Alzheimer's disease and frontotemporal dementia. *Curr Mol Med* **14**, 993-1000.
- [24] Zhan XH, Zha GC, Jiao JW, Yang LY, Zhan XF, Chen JT, Xie DD, Eyi UM, Matesa RA, Obono MM, Ehapo CS, Wei EJ, Zheng YZ, Yang H, Lin M (2015) Rapid identification of apolipoprotein E genotypes by high-resolution melting analysis in Chinese Han and African Fang populations. *Exp Ther Med* **9**, 469-475.
- [25] Duits FH, Teunissen CE, Bouwman FH, Visser PJ, Mattsson N, Zetterberg H, Blennow K, Hansson O, Minthon L, Andreassen N, Marcusson J, Wallin A, Rikkert MO, F, Grafman J, Cappa SF, Freedman R, Blalock EM, Bales KR, BL (2011) Sensitivity of revised diagnostic criteria for the behavioural variant of frontotemporal dementia. *Brain* **134**, 2456-2477.
- [26] Giaccone G, Rossi G, Morbin M, Tassi P, Ghisleni A, Bruni A (2002) A713T mutation of the AβPP gene in an Italian family with Alzheimer disease and frontotemporal angiopathy. *Neurobiol Aging* **23**(Suppl 1), 105-109.
- [27] Bernardi L, Geracitano S, Colao F, Anfosso M, Frangipane F, Curcio F, Tomaino C, Vasso F, Smirne N, et al. (2009) AβPP A713T mutation in a family with Alzheimer disease with cerebrovascular lesions. *Neurology* **72**, 383-389.
- [28] Guerreiro RJ, Baquero M, Blesa N, Bullido MJ, Calado A, Crook R, Gómez-Isla T, Hernández I, Lleó A, Lage P, Masdeu J, Molina-Porcel L, Pérez-Tur J, Relvas R, Oliveira CR, et al. (2004) The A713T mutation of the AβPP gene in a family with Alzheimer's disease yields novel mutations in presenilin 1. *Neurobiol Aging* **25**, 725-731.
- [29] Cruts M, Theuns J, Van Broeckhoven C, Blalock EM, Bales KR, BL (2004) Specific mutation databases for neurodegenerative diseases. *Hum Mutat* **23**, 1340-1344.
- [30] Kinoshita J, Clark T (2007) Alzheimer's disease and frontotemporal dementia. *Hum Mutat* **28**, 365-381.
- [31] Stenson PD, Mort M, Ball EV, Shapiro DN (2014) The Human Gene Mutation Database: a comprehensive mutation repository for clinical genetics, diagnostic testing and research. *Hum Genet* **133**, 1-9.
- [32] Schöll M, Wall A, Thordardottir S, N, Långström B, Almkvist O, Graff C, Low PiB PET retention in presenile AD: A study of biomarkers in Arctic APP mutation carriers. *Neurobiol Aging* **29**, 229-236.
- [33] Tomiyama T, Nagata T, Shimada H, A, Kanemitsu H, Takuma H, Kuwano M, Ataka S, Wada Y, Yoshioka E, Nishimori H (2008) A new amyloid variant in Alzheimer's-type dementia. *Neurobiol Aging* **29**, 377-387.

AUTHOR COPY